

# Glutamine Synthetases of Higher Plants<sup>1</sup>

EVIDENCE FOR A SPECIFIC ISOFORM CONTENT RELATED TO THEIR POSSIBLE PHYSIOLOGICAL ROLE AND THEIR COMPARTMENTATION WITHIN THE LEAF

Received for publication September 8, 1982 and in revised form December 11, 1982

SHEILA F. McNALLY, BERTRAND HIREL, PIERRE GADAL, A. FRED MANN, AND GEORGE R. STEWART  
School of Botany, Trinity College, University of Dublin, Dublin 2, Eire (S. F. McN.); Physiologie Végétale  
Métabolique, ERA au CNRS No. 799, Université de Paris Sud, Bat 430, 91405 Orsay Cedex, France (B. H., P.  
G.); and Department of Botany, Birbeck College, University of London, London WC1E 7HX, Great Britain (A.  
F. M., G. R. S.)

## ABSTRACT

The chromatographic properties of glutamine synthetase isoforms have been investigated in a wide range of higher plant leaves and shoots using ion exchange chromatography. Different patterns of glutamine synthetase isoform content were observed. Among higher plants, four patterns or groups could be recognized. The first group is characterized by having only cytosolic glutamine synthetase, whereas the second group is distinguished by having only chloroplastic glutamine synthetase. The third group is characterized by cytosolic glutamine synthetase being a minor component of the total leaf glutamine synthetase activity. The fourth group is distinct from the other groups in having high cytosolic and chloroplast glutamine synthetase activity. Immunological studies have been undertaken on a few species from each group to identify unambiguously both cytosolic and chloroplastic glutamine synthetases.

It is now established that glutamine synthesis may occur in two compartments of the leaf cells of most higher plants (1, 6-8, 15, 16, 27). Chromatographic and electrophoretic techniques combined with subcellular localization studies have demonstrated the presence of two different isoforms of GS in green tissue (1, 6-8, 13, 15, 16, 18, 24). One isoform, designated GS<sub>1</sub>, is localized in the cytosol, whereas the other, designated GS<sub>2</sub>, is found in the chloroplast (7-9, 15). The isoforms have been characterized by their specific kinetic and regulatory properties. They exhibit differences in their pH optima, heat stability, and  $K_m$  for glutamate (6, 13, 15).

Within the small number of plant species studied such as barley (15), rice (6), pea (7), sorghum (8), spinach (9), and pumpkin (13), differences have been observed in GS isoform complement. In spinach (9), only GS<sub>2</sub> is present, whereas in other C<sub>3</sub> plants like rice (6), pea (7), and barley (15), a minor GS<sub>1</sub> component is present (up to 20% of total enzyme activity). In sorghum, a C<sub>4</sub> plant, the pattern is yet again different with GS<sub>1</sub> representing about 70% of total GS activity in the leaf (8). We have extended these studies to a wider range of higher plants including halophytes and parasitic plants to confirm differences in the relative propor-

tions of GS<sub>1</sub> to GS<sub>2</sub>.

Immunological methods were also employed with selected plant species to identify very specifically the cytosolic and chloroplastic form of GS.

## MATERIALS AND METHODS

**Plant Culture.** Most plants were grown in plastic containers filled with vermiculite and watered daily for 3 weeks before harvesting, except for *Alnus glutinosa* which was grown for 6 months, with a complete Hoagland solution (11) in a controlled environment chamber (7). Tropical grasses were grown on a Lockard solution (14) under controlled conditions (6). CAM plants were grown for 8 weeks at the Phytotron of Gif-sur-Yvette (France) on a nutrient solution (4). Parasitic plant species were collected or grown as previously described (18). Halophytic plant species were collected from a salt marsh at Ichenore (West Sussex, England) or grown from seed in John Innes Compost No. 3 (1).

**Enzyme Extraction and Purification.** All operations were carried out at 4°C. Ten to 20 g of fresh leaves (groups B, C, and D) or shoots (group A) were used in each experiment. For most plants, the grinding medium (100 ml) contained 25 mM Tris-HCl buffer (pH 7.6), 1 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, 1 mM DTT, and 1 g Polyclar (acid-washed). Leaves or shoots were homogenized with a Polytron PT10 mixer at maximum speed for 4 × 15 s. The brei was filtered through two layers of gauze and then centrifuged at 45,000g for 30 min. The supernatant was directly layered on the top of a DEAE-Sephacel column (Pharmacia, Uppsala) (1 × 10 cm) previously equilibrated in the homogenizing buffer without Polyclar. A linear gradient of 0 to 400 mM NaCl or 0 to 600 mM KCl dissolved in 200 ml of the equilibrating buffer was used to elute the proteins. Four-ml fractions were collected and the flow rate adjusted to 20 ml h<sup>-1</sup>.

The purification procedure for the following plants was modified as follows.

*Alnus glutinosa.* The grinding medium was 50 mM Tris-HCl,

Table 1. Plants Exhibiting Only GS<sub>1</sub> Activity: Group A

| Plant Species                             |
|---|
| <i>Cuscuta australis</i> (P) <sup>a</sup> |
| <i>Lathraea clandestina</i> (P)           |
| <i>Lathraea squamosa</i> (P)              |
| <i>Orobancha cernua</i> (P)               |
| <i>Orobancha hederæ</i> (P)               |
| <i>Orobancha minor</i> (P)                |
| <i>Orobancha ramosa</i> (P)               |

<sup>a</sup> P, parasite.

<sup>1</sup> Supported by Grant Recherche Cooperative sur Programme no. 389, by Grant no. 7870445 for the Délégation Générale à la Recherche Scientifique et Technique, and by the Agricultural Research Council.

<sup>2</sup> Abbreviations: GS, glutamine synthetase; GS<sub>1</sub>, cytosolic glutamine synthetase; GS<sub>2</sub>, chloroplastic glutamine synthetase.




FIG. 1. Representative elution profiles of GS<sub>1</sub> and GS<sub>2</sub> of the four major groups of plants after chromatography on DEAE-Sephacel ion-exchange resin. Results are expressed as a percentage of total activity in the leaf. Recovery of GS activity after chromatography was always greater than 80% of the total activity initially loaded into the column. A, Group A. The achlorophyllous parasite *Cuscuta australis* (Table I). Maximum activity is 0.13 nkatal g<sup>-1</sup> fresh wt. B, Group B. The solanaceae *Lycopersicon esculentum* (Table II). Maximum activity is 0.66 nkatal g<sup>-1</sup> fresh wt. C, Group C. The legume *Phaseolus vulgaris* (Table III). Maximum activity is 0.45 nkatal g<sup>-1</sup> fresh wt. D, Group D. A tropical grass *Chloris gayana* (Table IV). Maximum activity is 0.58 nkatal g<sup>-1</sup> fresh wt.

Table II. Plants Exhibiting Only GS<sub>2</sub> Activity: Group B

| Plant Species                              |
|--|
| <i>Ammophila arenaria</i> (G) <sup>a</sup> |
| <i>Atriplex hastata</i> (H)                |
| <i>Atriplex lanceniata</i> (H)             |
| <i>Atriplex patula</i> (H)                 |
| <i>Cakile maritima</i> (H)                 |
| <i>Crithmum maritimum</i> (H)              |
| <i>Croixalacrma jobi</i>                   |
| <i>Croton megacarpus</i>                   |
| <i>Drepanocarpus lunatus</i>               |
| <i>Eriophorum angustifolium</i> (G)        |
| <i>Frankenia laevis</i> (H)                |
| <i>Halimione portulacoides</i> (H)         |
| <i>Lemna minor</i>                         |
| <i>Limonium vulgare</i> (H)                |
| <i>Lupinus</i> sp. (L)                     |
| <i>Lycopersicon esculentum</i>             |
| <i>Nicotiana tabacum</i>                   |
| <i>Petunia grandiflora</i>                 |
| <i>Salicornia europaea</i> (H)             |
| <i>Senecio vulgaris</i>                    |
| <i>Spinacia oleracea</i>                   |
| <i>Suaeda maritima</i> (H)                 |
| <i>Marchantia polymorpha</i> (LP)          |
| <i>Osmunda regalis</i> (LP)                |

<sup>a</sup> G, Graminae; H, halophyte; L, legume; LP, lower plant.

(pH 7.8), containing 2 mM MgCl<sub>2</sub>, 2 mM EDTA, 50 mM 2-mercaptoethanol, 1 mM DTT, 2% (w/v) soluble PVP (PVP40, Sigma), and 2 g Polyclar (acid-washed). After filtration and centrifugation, the brei was diluted twice with cold distilled H<sub>2</sub>O and directly applied onto a DEAE-Sephacel column.

*Kalanchoë blossfeldiana* and *Kalanchoë daigremontiana*. The

Table III. Plants Exhibiting GS<sub>1</sub> Activity Equal to or Lower than 30% of Total Activity: Group C

Relative proportions of GS<sub>1</sub> and GS<sub>2</sub> are expressed as a percentage of total activity in the leaf.

| Plant Species                               | GS <sub>1</sub> | GS <sub>2</sub> |
|---|-----------------|-----------------|
| %   |                 |                 |
| <i>Alnus glutinosa</i> (T) <sup>a</sup>     | 30              | 70              |
| <i>Aster tripolium</i> (H)                  | 27              | 63              |
| <i>Avena sativa</i> (G)                     | 10              | 90              |
| <i>Dendrobium nobile</i> (O)                | 22              | 78              |
| <i>Festuca rubra</i> (H)                    | 10              | 90              |
| <i>Hordeum vulgare</i> (G)                  | 17              | 83              |
| <i>Oryza sativa</i> (G)                     | 18              | 82              |
| <i>Panicum maximum</i> (C <sub>4</sub> PCK) | 15              | 85              |
| <i>Phaseolus aureus</i> (L)                 | 25              | 75              |
| <i>Phaseolus vulgaris</i> (L)               | 5               | 95              |
| <i>Pisum sativum</i> (L)                    | 15              | 85              |
| <i>Plantago maritima</i> (H)                | 9               | 91              |
| <i>Puccinellia maritima</i> (H)             | 20              | 80              |
| <i>Spergularia marina</i> (H)               | 15              | 85              |
| <i>Triticum aestivum</i> (G)                | 18              | 82              |

<sup>a</sup> T, tree; H, halophyte; G, Graminae; O, orchid; C<sub>4</sub> PCK, plant with a C<sub>4</sub> carbon metabolism pyruvate carboxykinase type; L, legume.

Table IV. Plants Exhibiting GS<sub>2</sub> Activity Greater than 30% of Total Activity: Group D

Relative proportions of GS<sub>1</sub> and GS<sub>2</sub> are expressed as a percentage of total activity in the leaf.

| Plant Species                                   | GS <sub>1</sub> | GS <sub>2</sub> |
|---|-----------------|-----------------|
| %   |                 |                 |
| <i>Chloris gayana</i> (C <sub>4</sub> PCK)      | 45              | 55              |
| <i>Digitaria sanguinalis</i> (C <sub>4</sub> )  | 80              | 20              |
| <i>Echinochloa crus-galli</i> (C <sub>4</sub> ) | 71              | 29              |
| <i>Erythrina abyssinica</i> (TL)                | 70              | 30              |
| <i>Erythrina cristagalli</i> (TL)               | 51              | 49              |
| <i>Erythrina corallodendron</i> (TL)            | 45              | 55              |
| <i>Erythrina flabelliforme</i> (TL)             | 68              | 32              |
| <i>Kalanchoë blossfeldiana</i> (CAM)            | 65              | 35              |
| <i>Kalanchoë daigremontiana</i> (CAM)           | 36              | 64              |
| <i>Pennisetum americanum</i> (C <sub>4</sub> )  | 50              | 50              |
| <i>Sorghum vulgare</i> (C <sub>4</sub> )        | 70              | 30              |
| <i>Zea mays</i> (C <sub>4</sub> )               | 45              | 55              |

<sup>a</sup> C<sub>4</sub> PCK, plant with a C<sub>4</sub> carbon metabolism pyruvate carboxykinase type; C<sub>4</sub>, plant with a C<sub>4</sub> carbon metabolism; TL, tropical legume.

grinding medium was 200 mM Tris-HCl (pH 8.0) containing 1 mM MgCl<sub>2</sub>, 1 mM EDTA, 10 mM 2-mercaptoethanol, 1 mM DTT, 2% (w/v) PEG 20,000 (Serva). After filtration and centrifugation, the brei was dialyzed against a large excess of 10 mM Tris-HCl (pH 7.6) containing 1 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, 1 mM DTT for 12 h and then applied to a DEAE-Sephacel column equilibrated in the same buffer.

**Halophytic Species.** Crude extracts were first desalted on G25 equilibrated with extraction buffer (minus Polyclar) before chromatography on DEAE-Sephacel).

**Enzyme Assays.** Determination of transferase and synthetase activities of GS were as described by Rhodes *et al.* (23).

**Immunodiffusion and Immunoprecipitation.** Immunodiffusion and immunoprecipitation experiments were performed as previously described (9).

## RESULTS AND DISCUSSION

Four groups of plants can be distinguished by their specific GS isoform content. The first group of plants (Table I, group A) is




FIG. 2. Immunochemical characterization of GS<sub>1</sub> and GS<sub>2</sub> from pea leaves. Immunodiffusion was performed on agar plates (2%) according to the Ouchterlony technique (21). Three wells on the left contained 10  $\mu$ l of concentrated GS<sub>2</sub> and three wells on the right contained 10  $\mu$ l of concentrated GS<sub>1</sub>. The central well contained 10  $\mu$ l of antiserum raised against GS from spinach chloroplasts (IgG). The plates were placed for 48 h at 4°C in a humid atmosphere and rinsed for 29 h in 0.9% NaCl solution. The plates were dried under filter paper at room temperature, stained for 1 h with Coomassie Brilliant Blue R, and destained overnight. Similar results were also obtained with GS<sub>1</sub> and GS<sub>2</sub> isolated from rice, sorghum, and *Kalanchoë* leaves.

distinct in having only one peak of GS activity eluting at about 0.15 M NaCl concentration (Fig. 1A). Similar elution values have been obtained for GS<sub>1</sub> from various plants (1, 6–8, 15, 16). This suggests that only GS<sub>1</sub> is present in this group of achlorophyllous higher plant parasites which are nutritionally dependent upon their hosts (19). The second group (Table II, group B) is characterized by the apparent absence of GS<sub>1</sub>, a pattern already described

for spinach (9). It can be seen in Figure 1B that a single peak of enzyme activity is present but which is eluted at a higher ionic strength of about 0.22 M NaCl. This elution profile is typical for example, of a number of chenopods from various environments, including salt marsh habitats. It is interesting to note that two lower plants, *Marchantia* and *Osmunda*, also possess only GS<sub>2</sub>. The third group (Table III, group C) is distinguished in having two GS isoforms, with a minor GS<sub>1</sub> (5–30% of total leaf enzyme activity) eluting at about 0.15 M NaCl and a major GS<sub>2</sub> (70–95% of total leaf enzyme activity) which is eluted at about 0.25 M NaCl (Fig. 1C). This pattern of GS activity is typical of many C<sub>3</sub> grasses and some temperate legumes. It is noteworthy that some halophytic plant species possess both GS isoforms, suggesting that GS isoform complement is not necessarily dependent upon environmental conditions. In the fourth group (Table IV, group D), GS<sub>1</sub> and GS<sub>2</sub> are also present, eluting at 0.15 and 0.25 M NaCl concentrations, respectively (Fig. 1D), but as shown in Table IV, the contribution of GS<sub>1</sub> is consistently higher than that found in Group C (Table III). In corn, for example, GS<sub>1</sub> activity represents about 45% of total enzyme activity in the leaf, and in *Digitaria* the contribution of GS<sub>1</sub> is much higher, up to 80% of total GS. This pattern is observed for most C<sub>4</sub> plants, a CAM plant, *Kalanchoë*, and some tropical legumes such as *Erythrina*.

Immunodiffusion and immunoprecipitation were employed to characterize unambiguously GS<sub>1</sub> and GS<sub>2</sub> isoforms from some of the plant species examined using antibodies raised against chloroplastic GS from spinach leaves (9). It can be seen in Figure 2 that these antibodies recognize the chloroplastic GS from pea leaves but also GS<sub>1</sub>, the cytosolic enzyme, with a faint band of precipitate. Furthermore, it can be observed that a partial identity exists between the two proteins with a spur occurring between the two precipitin bands. Similar results were also obtained, e.g. in rice, sorghum, and *Kalanchoë* which differ in their relative proportions of GS isoform content. The immunological behavior of GS<sub>1</sub> and GS<sub>2</sub> was also studied by immunoprecipitation. GS<sub>1</sub> and GS<sub>2</sub> were separated by ion-exchange chromatography on DEAE-Sephacel and immunoprecipitated. Constant amounts of enzyme are incubated with increasing concentrations of antibodies and the enzyme activity is measured in the supernatant following the removal of the immune complexes. Figure 3B shows the results obtained with rice; similar results were observed with GS<sub>1</sub> and GS<sub>2</sub> isolated from pea and sorghum leaves. For GS<sub>1</sub>, only 40% of total enzyme activity is precipitated with 200  $\mu$ l of antibodies. GS<sub>2</sub>, on the other hand, is completely precipitated with 100  $\mu$ l of




FIG. 3. Immunoprecipitation curves of GS from spinach (A) and rice (B) leaves. In B, GS<sub>1</sub> and GS<sub>2</sub> were separated by using DEAE-Sephacel chromatography and submitted separately to immunoprecipitation with antibodies raised against the spinach chloroplastic GS. Constant amounts of GS<sub>1</sub> and GS<sub>2</sub> activity (0.83 nkat) were incubated with increasing volume of either antiserum or nonimmune serum. Samples were incubated for 1 h at 37°C, then overnight at 4°C, and finally centrifuged at 10,000g for 15 min. GS activity was assayed in the supernatant. A, GS from spinach leaves (●). B, GS<sub>1</sub> from rice leaves (○); GS<sub>2</sub> from rice leaves (●). A and B, control with nonimmune serum (■).

antibodies. This latter result is comparable to that obtained for spinach (Fig. 3A). This result suggests that GS<sub>2</sub> isolated from a number of different species are similar. Moreover, this immunochemical approach shows that GS<sub>1</sub> and GS<sub>2</sub> in several vascular plants are different proteins. This confirms the differences already obtained with comparative studies of their kinetic and regulatory properties (6, 13, 15).

It has been suggested that GS<sub>1</sub> is responsible for the reassimilation of ammonia produced during photorespiration (6, 12, 15) and also in glutamine synthesis in the dark (6). GS<sub>2</sub> has been suggested to function mainly in primary ammonia assimilation in the chloroplast. This activity is probably a light-dependent process controlled by ATP, Mg<sup>2+</sup>, and H<sup>+</sup> concentrations in the stroma (6, 10). Very recently, it has been demonstrated that the chloroplast has the capacity to directly reassimilate ammonia released during photorespiration (2, 28). Furthermore, it has been demonstrated that the level of GS activity in the chloroplast is much higher than glutamate synthase (25) and could play a role in plastid ammonia detoxification in addition to its assimilatory function (5, 22). Clearly defined physiological roles of cytosolic and chloroplastic GS are more complicated than was first proposed in pea leaf cells (12, 27).

In several achlorophyllous plant parasites, only GS<sub>1</sub> is present Table I, (group A). This suggests that in these plants, glutamine synthesis can occur only in the cytosol which is comparable to the pattern observed in etiolated tissue where GS<sub>2</sub> activity is low (6, 7) or absent (8, 15). Moreover in nonphotosynthetic tissues such as roots, it has been shown that GS occurs in the cytosol (20, 26). These results suggest that GS<sub>1</sub> does not function exclusively in the reassimilation of ammonia released during the decarboxylation of glycine to serine but is also capable of primary ammonia assimilation. In contrast, some plants have only GS<sub>2</sub>, the chloroplastic isoform Table II, (group B) and this obviously implies that ammonia resulting from any deamination of glycine must be transported to the chloroplast for reassimilation. This result is in agreement with those already obtained for spinach leaves (2, 28) where it has been shown that the chloroplastic GS is present in sufficient quantities to allow the reassimilation of ammonia resulting both from photorespiration and from nitrate reduction.

In a recent paper (17), it has been estimated that the flux of ammonia through the photorespiratory nitrogen cycle can be 5 to 10 times higher than for nitrate reduction. Therefore, the pattern of GS activity exhibited by many C<sub>3</sub> plants which contain less than 30% of total GS activity as GS<sub>1</sub> activity Table III, (group C) does not allow for the reassimilation of all ammonia released during photorespiration by cytosolic GS as described by Keys *et al.* (12). Consequently, in these plants which have high rates of photorespiration it seems likely that GS<sub>2</sub> must function in the reassimilation of the ammonia released during the decarboxylation of glycine to serine.

The group of plants which most closely approach the Keys *et al.* model (12) are the C<sub>4</sub> plants (Table IV, group D), where GS<sub>1</sub> activity is present at higher levels. However, photorespiration in C<sub>4</sub> plants as measured by CO<sub>2</sub> evolution is considered to be very low or absent (3). These plants have ribulose-1,5-bisphosphate carboxylase similar to that of C<sub>3</sub> plants and have the necessary C<sub>2</sub> enzymes for glycine formation and deamination. The present results suggest that C<sub>4</sub> plants may reassimilate photorespiratory ammonia via cytosolic GS. Furthermore, GS activity on a fresh weight basis is very similar between C<sub>3</sub> and C<sub>4</sub> plants. This suggests that GS<sub>2</sub> must play a role in the reassimilation of ammonia arising from the photorespiratory nitrogen cycle in the majority of C<sub>3</sub> plants examined.

In the results presented here, four groups of plants displaying

different GS isoform complements could be distinguished, characterized mainly by the increasing importance of GS<sub>1</sub>. Clarification of the possible physiological roles of GS<sub>1</sub> and GS<sub>2</sub> suggested in the present paper will require further experimentation, particularly with influence of light intensity, daylength on C<sub>4</sub>, and tropical plant nitrogen metabolism. It also seems worthwhile to determine the differential gene expression of GS<sub>1</sub> and GS<sub>2</sub> among the four different groups of plants.

#### LITERATURE CITED

1. AHMAD I, F LARHER, AF MANN, SF McNALLY, GR STEWART 1982 Nitrogen metabolism of halophytes. IV. Characteristics of glutamine synthetase from *Triglochin maritima* L. New Phytol 91: 585-595
2. BERGMAN A, P GARDESTROM, I ERICSON 1981 Release and refixation of ammonia during photorespiration. Physiol Plant 53: 528-537
3. BLACK CC 1973 Photosynthetic carbon fixation in relation to net CO<sub>2</sub> uptake. Annu Rev Plant Physiol 24: 253-286
4. CHOUARD P, R JACQUES, N DE BILDERLING 1973 Phytotron Phytotronique. Endeavour 31: 41-45
5. GIVAN CV 1979 Metabolic detoxification of ammonia in higher plants. Phytochemistry 18: 375-382
6. HIREL B, P GADAL 1980 Glutamine synthetase in rice. A comparative study of the enzymes from roots and leaves. Plant Physiol 66: 619-623
7. HIREL B, P GADAL 1981 Glutamine synthetase isoforms in pea leaves. Intracellular localization. Z Pflanzenphysiol 102: 315-319
8. HIREL B, P GADAL 1982 Glutamine synthetase in a C<sub>4</sub> plant: *Sorghum vulgare* L. Physiol Plant 54: 69-74
9. HIREL B, C PERROT-RECHENMANN, A SUZUKI, J VIDAL, P GADAL 1982 Glutamine synthetase in spinach leaves. Immunological studies and immuno-cytochemical localization. Plant Physiol 69: 983-987
10. HIREL B, J VIDAL, P GADAL 1982 Evidence for a cytosolic dependent light induction of chloroplastic glutamine synthetase in etiolated rice leaves. Planta 155: 17-23
11. HOAGLAND DR, DI ARNON 1938 The water culture for growing plants without soil. Circ Calif Agric Exp Stn 347: 461-462
12. KEYS AJ, IF BIRD, MF CORNELIUS, PJ LEA, RM WALLSGROVE, BJ MIFLIN 1978 Photorespiratory nitrogen cycle. Nature 275: 741-742
13. KRETOVICH WL, ZG EVSTIGNEVA, AV PUSHKIN, TZ DZOKHARIDZE 1981 Two forms of glutamine synthetase in leaves of *Cucurbita pepo*. Phytochemistry 20: 625-629
14. LOCKARD RG 1959 In Sand and Water Culture Methods. 4. Hewitt Eds, Table 30c, pp 547
15. MANN AF, PA FENTEM, GR STEWART 1979 Identification of two forms of glutamine synthetase in barley (*Hordeum vulgare* L.) Biochem Biophys Res Commun 88: 515-521
16. MANN AF, PA FENTEM, GR STEWART 1980 Tissue localization of barley (*Hordeum vulgare* L.) glutamine synthetase isoenzymes. FEBS Lett 110: 265-267
17. MIFLIN BJ, PJ LEA, RM WALLSGROVE 1980 The role of glutamine in ammonia assimilation and reassimilation in plants. In J. Mora, R. Palacios, eds, Glutamine: Metabolism, Enzymology and Regulation. Academic Press, New York, pp 213-234
18. McNALLY SF, GR STEWART 1982 Modifications in the N-metabolism of *Striga* associated with its parasitic habit. Plant Physiol 69: S-110
19. MUSSELMAN 1980 The biology of *Striga*, *Orobancha* and other root parasitic weeds. Annu Rev Phytopathol 18: 463-489
20. OAKS A, P GADAL 1979 Nitrate utilization in cells of higher plants. In C Nover, F Lynen, M Mothes, eds, Cell Compartmentations Metabolic Channeling. VEB Gustav, Fisher-Verlag, Jena and Elsevier/North Holland Biomedical Press, Amsterdam, pp 245-254
21. OUCHTERLONY O 1949 Antigen-antibody reactions in gels. Acta Pathol Microbiol Scand 26: 507-515
22. PLATT SG, G ANTHON 1981 Ammonia accumulation and inhibition of photosynthesis in methionine sulfoximine-treated spinach. Plant Physiol 67: 509-513
23. RHODES D, GA RENDON, GR STEWART 1975 The control of glutamine synthetase level in *Lemna minor* L. Planta 125: 203-210
24. STASIEWICZ S, VL DUNHAM 1979 Isolation and characterization of two forms of GS from soybean hypocotyls. Biochem Biophys Res Commun 87: 627-634
25. SUZUKI A, P GADAL 1982 Glutamate synthase from rice leaves. Plant Physiol 69: 848-852
26. SUZUKI A, P GADAL, A OAKS 1981 Intracellular distribution of enzymes associated with nitrogen assimilation in roots. Planta 151: 457-461
27. WALLSGROVE RM, PJ LEA, BJ MIFLIN 1979 Distribution of the enzymes of nitrogen assimilation within the pea leaf cell. Plant Physiol 63: 232-236
28. WOO KC, BC OSMOND 1982 Stimulation of ammonia and 2-oxoglutarate-dependent O<sub>2</sub> evolution in isolated chloroplasts by dicarboxylate and the role of the chloroplast in photorespiratory nitrogen recycling. Plant Physiol 69: 591-596